

INJ-COL-2026 02

**Columvi®**  
Glofitamab



**1. DESCRIPTION**  
**1.1 THERAPEUTIC / PHARMACOLOGIC CLASS OF DRUG**  
Antineoplastic agent, monoclonal antibody (recombinant humanized immunoglobulin G1)  
ATC code: L01FX28

Pharmacotherapeutic group: Antineoplastic agents, other monoclonal antibodies and antibody drug conjugates

**1.2 TYPE OF DOSAGE FORM**  
Concentrate for solution for Infusion

**1.3 ROUTE OF ADMINISTRATION**  
Intravenous (IV) Infusion

**1.4 STERILE / RADIOACTIVE STATEMENT**  
Sterile Product

**1.5 QUALITATIVE AND QUANTITATIVE COMPOSITION**  
Active ingredient: glofitamab

Columvi is a preservative-free, colorless, clear solution supplied in colorless type I borosilicate glass single-dose vials with fluororesin-laminated rubber stopper and aluminum seal with plastic flip-off cap containing:

- 2.5 mg of glofitamab/2.5 mL at a concentration of 1 mg/mL
- 10 mg of glofitamab/10 mL at a concentration of 1 mg/mL

Excipients: D-Sucrose; L-Histidine; L-Histidine Hydrochloride, Monohydrate; L-Methionine; Polysorbate 20; Water for Injection.

**2. CLINICAL PARTICULARS**  
**2.1 THERAPEUTIC INDICATION(S)**

Columvi in combination with gemcitabine and oxaliplatin is indicated for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma not otherwise specified (DLBCL NOS) who are not candidates for autologous stem cell transplant (ASCT).

Columvi is indicated for the treatment of adult patients with relapsed or refractory diffuse large B-cell lymphoma (DLBCL) after two or more lines of systemic therapy. (See 3.1.2 Clinical / Efficacy Studies)

**2.2 DOSAGE AND ADMINISTRATION**

**General**  
Columvi therapy should only be administered under the supervision of a healthcare professional experienced in the treatment of cancer patients and who has access to appropriate medical support to manage severe reactions associated with cytokine release syndrome (CRS). At least 1 dose of tocilizumab for use in the event of CRS must be available prior to Columvi infusion at Cycles 1 and 2. Access to an additional dose of tocilizumab within 8 hours of use of the previous tocilizumab dose must be ensured. See Section 2.4.1 Warnings and Precautions.

**Pre-treatment with Obinutuzumab**

All patients must receive a single 1000 mg dose of obinutuzumab on Cycle 1 Day 1 (7 days prior to initiation of Columvi treatment); see Table 2, Table 3 and *Delayed or Missed Doses*. This is to deplete circulating and lymphoid tissue B cells and thereby reduce the risk of CRS.

Obinutuzumab should be administered as an intravenous infusion at 50 mg/h. The rate of infusion can be escalated in 50 mg/h increments every 30 minutes to a maximum of 400 mg/h.

Refer to the obinutuzumab prescribing information for complete information on premedication, preparation, administration, and management of adverse reactions of obinutuzumab.

**Premedication and Prophylactic Medications**

**Cytokine release syndrome prophylaxis**

Columvi should be administered to well-hydrated patients. Premedication to reduce the risk of CRS (see Section 2.4.1 Warnings and Precautions) is outlined in Table 1.

**Table 1 Premedication Before Columvi Infusion To Reduce the Risk of Cytokine Release Syndrome**

Treatment Cycle (Day)	Patients requiring premedication	Premedication	Administration
Cycle 1 (Day 8, Day 15); Cycle 2 (Day 1); Cycle 3 (Day 1)	All patients	20 mg intravenous dexamethasone <sup>a</sup>	Completed at least 1 hour prior to Columvi infusion.
		Oral analgesic / anti-pyretic <sup>b</sup> Anti-histamine <sup>c</sup>	At least 30 minutes before Columvi infusion.
All subsequent infusions	All patients	Oral analgesic / anti-pyretic <sup>b</sup> Anti-histamine <sup>c</sup>	At least 30 minutes before Columvi infusion.
	Patients who experienced CRS with previous dose	20 mg intravenous dexamethasone <sup>a</sup>	Completed at least 1 hour prior to Columvi infusion.

a. If patient has an intolerance to dexamethasone or dexamethasone is unavailable, administer 100 mg prednisone/prednisolone or 80 mg methylprednisolone.  
b. For example, 1000 mg acetaminophen/paracetamol.  
c. For example, 50 mg diphenhydramine.

**Infection prophylaxis**

Prophylaxis is recommended to reduce the risk of infection (see Section 2.4 Warnings and Precautions).

Consider prophylaxis for cytomegalovirus (CMV), herpes, pneumocystis jirovecii pneumonia, and other opportunistic infections in patients at increased risk (see Section 2.6.1 Description of selected adverse reactions).

**Recommended Dosage**

Columvi dosing begins with a step-up dosing schedule (which is designed to decrease the risk of CRS), leading to the recommended dose of 30 mg.

**Columvi Monotherapy Dose Step-Up Schedule**

Columvi must be administered as an intravenous infusion according to the dose step-up schedule leading to the recommended dosage of 30 mg (as shown in Table 2), after completion of pre-treatment with obinutuzumab on Cycle 1 Day 1. Each cycle is 21 days.

**Table 2 Columvi Monotherapy Dose Step-Up Schedule for Patients with Relapsed or Refractory DLBCL**

Treatment Cycle, Day <sup>a</sup>	Dose of Columvi	Duration of infusion
Cycle 1 (Pre-treatment and step-up dose)	Day 1	Pre-treatment with obinutuzumab 1000mg <sup>b</sup>
	Day 8	2.5 mg
	Day 15	10 mg
Cycle 2	Day 1	30 mg
Cycle 3 to 12	Day 1	30 mg

a. Each treatment cycle is 21 days.  
b. Refer to *Pre-treatment with obinutuzumab* described above.  
c. For patients who experience CRS with their previous dose of Columvi, the duration of infusion may be extended up to 8 hours (see Table 4 and Section 2.4.1 Warnings and Precautions).

d. At the discretion of the treating physician, if the previous infusion was well tolerated. If the patient experienced CRS with the previous dose, the duration of infusion should be maintained at 4 hours.

**Columvi Dose Step-up Schedule in Combination with Gemcitabine and Oxaliplatin**

Columvi must be administered as an intravenous infusion according to the dose step-up schedule leading to the recommended dosage of 30 mg (as shown in Table 3), after completion of pre-treatment with obinutuzumab on Cycle 1 Day 1. Administer Columvi in combination with gemcitabine and oxaliplatin at Cycles 1-8 and as monotherapy at Cycles 9-12. Each cycle is 21 days.

**Table 3 Columvi Dose Step-Up Schedule in Combination with Gemcitabine and Oxaliplatin for Patients with Relapsed or Refractory DLBCL**

Treatment Cycle, Day <sup>a</sup>	Dose of Columvi (duration of infusion)	Dose of gemcitabine	Dose of oxaliplatin
Cycle 1 (pre-treatment and step-up dose)	Day 1	Pre-treatment with obinutuzumab 1000 mg <sup>b</sup>	
	Day 2	--	1000 mg/m <sup>2</sup> c
	Day 8	2.5 mg (4 hours) <sup>d</sup>	--
Cycle 2	Day 1	10 mg (4 hours) <sup>d</sup>	--
	Day 15	30 mg (4 hours) <sup>d,e</sup>	1000 mg/m <sup>2</sup> c
Cycle 3 to 8	Day 1	30 mg (2 hours) <sup>e,f</sup>	1000 mg/m <sup>2</sup> c
	Day 1	30 mg (2 hours) <sup>f</sup>	N/A

N/A=not applicable.  
a. Each treatment cycle is 21 days.  
b. Refer to *Pre-treatment with obinutuzumab* described above.  
c. Cycles 1-8: administer gemcitabine before oxaliplatin.  
d. For patients who experience CRS with their previous dose of Columvi, the duration of infusion may be extended up to 8 hours (see Table 4 and Section 2.4.1 Warnings and Precautions).  
e. Cycles 2-8: administer Columvi before gemcitabine and oxaliplatin. Gemcitabine and oxaliplatin may be given on Day 1 or 2.  
f. Infusion time may be shortened to 2 hours at the discretion of the treating physician, if the previous infusion was well tolerated. If the patient experienced CRS with the previous dose, the duration of infusion should be maintained at 4 hours.

**Monitoring after infusion**

- When Columvi is given as monotherapy, patients must be monitored for signs and symptoms of potential CRS during infusion and for at least 10 hours after completion of the infusion of the first Columvi dose (2.5 mg on Cycle 1 Day 8).
- When Columvi is given in combination with gemcitabine and oxaliplatin, patients must be monitored for signs and symptoms of potential CRS during all Columvi infusions and for 4 hours after completion of infusion of the first Columvi dose (2.5 mg on Cycle 1 Day 8).
- Patients who experienced Grade ≥ 2 CRS with their previous infusion should be monitored after completion of the infusion. See Table 4.
- Refer to section 2.6.1. *Description of selected adverse drug reactions from clinical trials, Cytokine Release Syndrome.*

All patients must be counselled on the risk, signs, and symptoms of CRS and advised to contact the healthcare provider immediately should they experience signs and symptoms of CRS.

**Duration of Treatment**

Treatment with Columvi monotherapy is recommended for a maximum of 12 cycles or until disease progression or unmanageable toxicity, whichever occurs first.

Treatment with Columvi in combination with gemcitabine and oxaliplatin is recommended for 8 cycles, followed by 4 cycles of Columvi monotherapy for a maximum of 12 cycles of Columvi in total or until disease progression or unmanageable toxicity, whichever occurs first.

**Delayed or Missed Doses**

**During step-up dosing (weekly dosing):**

- Following pre-treatment with obinutuzumab, if the Columvi 2.5 mg dose is delayed by more than 1 week, then repeat pre-treatment with obinutuzumab.
- Following Columvi 2.5 mg dose, if there is a Columvi treatment-free interval of 2 to 4 weeks, then repeat glofitamab 2.5 mg dose and resume the planned step-up dosing.
- Following Columvi 2.5mg dose, if there is a Columvi treatment-free interval of more than 4 weeks, then repeat pre-treatment with obinutuzumab and Columvi step-up dosing (see Cycle 1 in Table 2 and Table 3).
- Following Columvi 10 mg dose, if there is a Columvi treatment-free interval of 2 weeks to 6 weeks, then repeat the last tolerated Columvi dose and resume the planned step-up dosing.
- Following Columvi 10 mg dose, if there is a Columvi treatment-free interval of more than 6 weeks, then repeat pre-treatment with obinutuzumab and Columvi step-up dosing (see Cycle 1 in Table 2 and Table 3).

**After Cycle 2 (30 mg dose):**

- If there is a Columvi treatment-free interval of more than 6 weeks between cycles, then repeat pre-treatment with obinutuzumab and Columvi step-up dosing (see Cycle 1 in Table 2 and Table 3), and then resume the planned treatment cycle (30 mg dose).

**Preparation and Administration of Columvi**

**Preparation**  
Columvi must be diluted by a healthcare professional using aseptic technique, prior to intravenous administration. See Section 4.2 *Special Instructions for Use, Handling, and Disposal.*

**Administration**

- Columvi must be administered as an intravenous infusion through a dedicated infusion line.
- Columvi must not be administered as an intravenous push or bolus.
- Columvi must not be mixed with other drugs.

**Dose Modifications**

No dose reductions of Columvi are recommended.

**Management of Cytokine Release Syndrome**

Cytokine release syndrome should be identified based on the clinical presentation (see Section 2.4 Warnings and Precautions). Patients should be evaluated for other causes of fever, hypoxia, and hypotension, such as infections or sepsis. If CRS is suspected, it should be managed according to the CRS management recommendations based on American Society for Transplantation and Cellular Therapy [ASTCT] consensus grading in Table 4.

**Table 4 ASTCT CRS Grading and CRS Management Guidance**

Grade <sup>a</sup>	CRS Management	For Next Scheduled Columvi Infusion
Grade 1 Fever ≥38 °C	If CRS occurs during infusion: <ul style="list-style-type: none"> <li>• Interrupt infusion and treat symptoms</li> <li>• Restart infusion at slower rate when symptoms resolve</li> <li>• If symptoms recur, discontinue current infusion</li> </ul>	<ul style="list-style-type: none"> <li>• Ensure symptoms are resolved for at least 72 hours prior to next infusion</li> <li>• Consider slower infusion rate<sup>b</sup></li> </ul>
	If CRS occurs post-infusion: <ul style="list-style-type: none"> <li>• Treat symptoms</li> </ul>	
Grade 2 Fever ≥38 °C and/or hypotension not requiring vasopressors	If CRS lasts more than 48h after symptomatic management: <ul style="list-style-type: none"> <li>• Consider corticosteroids<sup>c</sup></li> <li>• Consider tocilizumab<sup>d</sup></li> </ul>	
	If CRS occurs during infusion: <ul style="list-style-type: none"> <li>• Discontinue current infusion and treat symptoms</li> <li>• Administer corticosteroids<sup>c</sup></li> <li>• Consider tocilizumab<sup>d</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Ensure symptoms are resolved for at least 72 hours prior to next infusion</li> <li>• Consider slower infusion rate<sup>b</sup></li> </ul>

and/or hypoxia requiring low-flow oxygen by nasal cannula or blow-by	If CRS occurs post-infusion: <ul style="list-style-type: none"> <li>• Treat symptoms</li> <li>• Administer corticosteroids<sup>c</sup></li> <li>• Consider tocilizumab<sup>d</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Monitor patients post-infusion</li> </ul>
--	---	--

**For Grade 2: Tocilizumab use**  
Do not exceed 3 doses of tocilizumab<sup>d</sup> in a period of 6 weeks.

If no prior use of tocilizumab or if 1 dose of tocilizumab was used within the last 6 weeks:

- Administer first dose of tocilizumab<sup>d</sup>
- If no improvement within 8 hours administer second dose of tocilizumab<sup>d</sup>
- After 2 doses of tocilizumab, consider alternative anti-cytokine and/or alternative immunosuppressant therapy

If 2 doses of tocilizumab were used within the last 6 weeks:

- Administer only one dose of tocilizumab
- If no improvement within 8 hours consider alternative anti-cytokine and/or alternative immunosuppressant therapy

Grade 3 Fever ≥38 °C and/or hypotension requiring a vasopressor (with or without vasopressin) and/or hypoxia requiring high-flow oxygen by nasal cannula, face mask, non-rebreather mask, or Venturi mask	If CRS occurs during infusion: <ul style="list-style-type: none"> <li>• Discontinue current infusion and treat symptoms</li> <li>• Administer corticosteroids<sup>c</sup></li> <li>• Administer tocilizumab<sup>d</sup></li> </ul>	<ul style="list-style-type: none"> <li>• Ensure symptoms are resolved for at least 72 hours prior to next infusion</li> <li>• Consider slower infusion rate<sup>b</sup></li> <li>• Monitor patients post-infusion</li> <li>• If Grade ≥ 3 CRS recurs at subsequent infusion, stop infusion immediately and permanently discontinue Columvi</li> </ul>
	If CRS occurs post-infusion: <ul style="list-style-type: none"> <li>• Treat symptoms</li> <li>• Administer corticosteroids<sup>c</sup></li> <li>• Administer tocilizumab<sup>d</sup></li> </ul>	

Grade 4 Fever ≥38 °C and/or hypotension requiring multiple vasopressors (excluding vasopressin) and/or hypoxia requiring oxygen by positive pressure (e.g., CPAP, BiPAP, intubation, and mechanical ventilation)	If CRS occurs during infusion or post-infusion: <ul style="list-style-type: none"> <li>• Permanently discontinue Columvi and treat symptoms</li> <li>• Administer corticosteroids<sup>c</sup></li> <li>• Administer tocilizumab<sup>d</sup></li> </ul>	

**For Grade 3 and Grade 4: Tocilizumab use**  
Do not exceed 3 doses of tocilizumab<sup>d</sup> in a period of 6 weeks.

If no prior use of tocilizumab or if 1 dose of tocilizumab was used within the last 6 weeks:

- Administer first dose of tocilizumab<sup>d</sup>
- If no improvement within 8 hours or rapid progression of CRS, administer second dose of tocilizumab<sup>d</sup>
- After 2 doses of tocilizumab, consider alternative anti-cytokine and/or alternative immunosuppressant therapy

If 2 doses of tocilizumab were used within the last 6 weeks:

- Administer only one dose of tocilizumab
- If no improvement within 8 hours or rapid progression of CRS, consider alternative anti-cytokine and/or alternative immunosuppressant therapy

- a. American Society for Transplantation and Cellular Therapy (ASTCT) 2019 consensus grading criteria.
- b. Duration of infusion may be extended up to 8 hours, as appropriate for that cycle (see Table 2).
- c. Corticosteroids (e.g., 10 mg IV dexamethasone, 100 mg IV prednisolone, 1-2 mg/kg IV methylprednisolone per day, or equivalent).
- d. Tocilizumab 8 mg/kg IV (not to exceed 800 mg).

**Management of Neurologic Toxicity, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS)**

Management recommendations for neurologic toxicity, including ICANS, are summarized in Table 5. At the first sign of neurologic toxicity, including ICANS, consider neurology evaluation and withholding Columvi based on the type and severity of neurotoxicity. Rule out other causes of neurologic symptoms. Provide supportive therapy, which may include intensive care.

**Table 5 Recommended Dosage Modification for Neurologic Toxicity (including ICANS)**

Adverse Reaction	Severity <sup>a,b</sup>	Actions
Neurologic Toxicity <sup>a</sup> (including ICANS <sup>b</sup> )  (see Section 2.4 Warnings and Precautions)	Grade 1	<ul style="list-style-type: none"> <li>• Continue Columvi and monitor neurologic toxicity symptoms.</li> <li>• If Grade 1 ICANS,<sup>b</sup> consider a single dose of dexamethasone 10 mg, if not taking other corticosteroids.</li> </ul>
	Grade 2	<ul style="list-style-type: none"> <li>• Withhold Columvi until neurologic toxicity symptoms improve to Grade 1 or baseline.<sup>c,d</sup></li> <li>• Provide supportive therapy and consider neurologic consultation and evaluation.</li> <li>• If Grade 2 ICANS,<sup>b</sup> treat with dexamethasone 10 mg intravenously every 12 hours, if not taking other corticosteroids, until improvement to Grade 1, then taper.</li> </ul>
	Grade 3	<ul style="list-style-type: none"> <li>• Withhold Columvi until neurologic toxicity symptoms improve to Grade 1 or baseline for at least 7 days.<sup>d,e</sup></li> <li>• For Grade 3 neurologic events lasting more than 7 days, consider permanently discontinuing Columvi.</li> <li>• Provide supportive therapy, which may include intensive care, and consider neurologic consultation and evaluation.</li> <li>• If Grade 3 ICANS,<sup>b</sup> treat with dexamethasone 10 mg intravenously every 6 hours, if not taking other corticosteroids, until improvement to Grade 1, then taper. Consider non-sedating anti-seizure medication for seizure prophylaxis until resolution of ICANS. Use anti-seizure medication for seizure management as needed.</li> </ul>
	Grade 4	<ul style="list-style-type: none"> <li>• Permanently discontinue Columvi.</li> <li>• Provide supportive therapy, which may include intensive care, and consider neurology consultation and evaluation.</li> <li>• If Grade 4 ICANS,<sup>b</sup> treat with dexamethasone 10 mg intravenously every 6 hours, if not taking other corticosteroids, until improvement to</li> </ul>

		Grade 1, then taper. Consider non-sedating anti-seizure medication for seizure prophylaxis until resolution of ICANS. Use anti-seizure medication for seizure management as needed.
--	--	---

a. Based on National Cancer Institute Common Terminology Criteria for Adverse Events (NCI CTCAE), version 4.03.

b. Based on ASTCT 2019 grading for ICANS.

c. Consider the type of neurologic toxicity before deciding to withhold Columvi.

d. See *Delayed or Missed doses section* on restarting Columvi after dose delays.

e. Evaluate benefit-risk before restarting Columvi.

## 2.2.1 Special Dosage Instructions

### Pediatric use

The safety and efficacy of Columvi in pediatric patients have not been established.

### Geriatric use

No dose adjustment of Columvi is required in patients  $\geq 65$  years of age. (See Section 2.5 *Use in Special Populations* and Section 3.2.5 *Pharmacokinetics in Special Populations*.)

### Renal Impairment

No dose adjustment of Columvi is required in patients with mild or moderate renal impairment (CrCL 30 to  $< 90$  mL/min). Columvi has not been studied in patients with severe renal impairment. (See Section 2.5 *Use in Special Populations* and Section 3.2.5 *Pharmacokinetics in Special Populations*.)

### Hepatic Impairment

No dose adjustment is required in patients with mild hepatic impairment (total bilirubin  $>$  upper limit of normal [ULN] to  $\leq 1.5 \times$  ULN or aspartate transaminase [AST]  $>$  ULN). No specific studies in patients with moderate or severe hepatic impairment have been conducted with Columvi. (See Section 2.5 *Use in Special Populations* and Section 3.2.5 *Pharmacokinetics in Special Populations*.)

## 2.3 CONTRAINDICATIONS

Columvi is contraindicated in patients with a known hypersensitivity to glofitamab or any of the excipients.

Refer to obinutuzumab-specific contraindications in the obinutuzumab prescribing information.

## 2.4 WARNINGS AND PRECAUTIONS

### 2.4.1 General

Refer to obinutuzumab-specific warnings and precautions in the obinutuzumab prescribing information.

In order to improve traceability of biological medicinal products, the trade name and the batch number of the administered product should be clearly recorded (or stated) in the patient file.

### CD20-negative disease

There are limited data available on patients with CD20-negative DLBCL treated with Columvi, and it is possible that patients with CD20-negative DLBCL may have less benefit compared to patients with CD20-positive DLBCL. The potential risks and benefits associated with treatment of patients with CD20-negative DLBCL with Columvi should be considered.

### Cytokine Release Syndrome

CRS, including life-threatening reactions, has been reported in patients receiving Columvi.

The most common manifestations of CRS were pyrexia, tachycardia, hypotension, chills, and hypoxia. Infusion-related reactions may be clinically indistinguishable from manifestations of CRS.

To reduce the occurrence of CRS, patients must be pre-treated with obinutuzumab, 7 days prior to initiation of Columvi, and should be premedicated with an antipyretic, anti-histamine, and a glucocorticoid. See Section 2.2 *Dosage and Administration*.

At least 1 dose of tocilizumab for use in the event of CRS must be available prior to Columvi infusion at Cycles 1 and 2. Access to an additional dose of tocilizumab within 8 hours of use of the previous tocilizumab dose must be ensured.

When Columvi is given as monotherapy, patients must be monitored during all Columvi infusions and for at least 10 hours after completion of the first infusion.

When Columvi is given in combination with gemcitabine and oxaliplatin, patients must be monitored during all Columvi infusions and for 4 hours after completion of the first infusion.

For complete information on monitoring, see Section 2.2 *Dosage and Administration*. The prescriber must counsel patients to seek immediate medical attention should signs or symptoms of CRS occur at any time.

Patients should be evaluated for other causes of fever, hypoxia, and hypotension, such as infections or sepsis. CRS should be managed based on the patient's clinical presentation and according to the CRS management guidance provided in Table 4 (see Section 2.2 *Dosage and Administration*).

Elevated liver function tests (AST and alanine transaminase [ALT]  $> 3 \times$  ULN and/or total bilirubin  $> 2 \times$  ULN) concurrent with CRS have been reported after Columvi use.

### Immunisation

The safety of immunisation with live vaccines during or following Columvi therapy has not been studied. Immunisation with live vaccines is not recommended during Columvi therapy.

### Neurologic Toxicity

Columvi can cause serious and fatal neurologic toxicity, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) (see Section 2.6.1 *Description of selected adverse events*). Grade 3 or higher neurologic adverse reactions occurred in 2.6% of patients. Cases of ICANS of any grade occurred in 7.1% of patients (see Section 2.6.1 *Description of selected adverse events*).

Co-administration of Columvi with other products that cause dizziness or mental status changes may increase the risk of neurologic toxicity. Optimize concomitant medications and hydration to avoid dizziness or mental status changes. Institute fall precautions as appropriate.

Monitor patients for signs and symptoms of neurologic toxicity, evaluate, and provide supportive therapy; withhold or permanently discontinue Columvi based on severity (see Section 2.2 *Dosage and Administration*).

Evaluate patients who experience neurologic toxicity such as tremors, dizziness, or adverse reactions that may impair cognition or consciousness promptly, including potential neurology evaluation. Advise affected patients to refrain from driving and/or engaging in hazardous occupations or activities, such as operating heavy or potentially dangerous machinery, until the neurologic toxicity fully resolves.

### Serious Infections

Serious infections, including opportunistic infections, have occurred in patients treated with Columvi (see Section 2.6.1 *Description of selected adverse events*).

Columvi must not be administered to patients with an active infection. Caution should be exercised when considering the use of Columvi in patients with a history of chronic or recurrent infection, those with underlying conditions that may predispose them to infections, or those who have had significant prior immunosuppressive treatment. Administer prophylactic antimicrobials, as appropriate. Patients should be monitored before and during Columvi treatment for the emergence of possible bacterial, fungal, and new or reactivated viral infections and treated appropriately.

Columvi should be temporarily withheld in the presence of an active infection until the infection has resolved. Patients should be instructed to seek medical advice if signs and symptoms suggestive of an infection occur.

Febrile neutropenia has been reported during treatment with Columvi. Patients with febrile neutropenia should be evaluated for infection and treated promptly.

### Tumor Flare

Tumor flare has been reported in patients receiving Columvi. Manifestations included localized pain and swelling (see Section 2.6.1 *Description of selected adverse events*).

Consistent with the mechanism of action of Columvi, tumor flare is likely due to the influx of T cells into tumor sites following Columvi administration and may mimic progression of disease. Tumor flare does not imply treatment failure or represent tumor progression.

Specific risk factors for tumor flare have not been identified, however, there is a heightened risk of compromise and morbidity due to mass effect secondary to tumor flare in patients with bulky tumors located in close proximity to airways and/or a vital organ. Monitoring and evaluation of tumor flare at critical anatomical sites is recommended in patients treated with Columvi and managed as clinically indicated.

### Tumor Lysis Syndrome

Tumor lysis syndrome (TLS) has been reported in patients receiving Columvi (see Section 2.6.1 *Description of selected adverse events*). Patients with high tumor burden, rapidly proliferative tumors, renal dysfunction, or dehydration are at greater risk of TLS.

Patients at risk should be monitored closely by appropriate clinical and laboratory tests for electrolyte status, hydration, and renal function. Appropriate prophylactic measures with anti-hyperuricemics (e.g., allopurinol or rasburicase) and adequate hydration should be considered prior to Columvi infusion.

Management of TLS may include aggressive hydration, correction of electrolyte abnormalities, anti-hyperuricemic therapy, and supportive care.

## 2.4.2 Drug Abuse and Dependence

Columvi does not have the potential for abuse and dependence.

## 2.4.3 Ability to Drive and Use Machines

Columvi has no or negligible influence on the ability to drive and use machines. Patients experiencing symptoms of CRS (pyrexia, tachycardia, hypotension, chills, hypoxia) should be advised not to drive or use machines until symptoms resolve.

## 2.5 USE IN SPECIAL POPULATIONS

### 2.5.1 Females and Males of Reproductive Potential

#### Fertility

See Section 3.3.3 *Impairment of fertility*.

#### Contraception

Female patients of reproductive potential must use highly effective contraceptive methods during treatment and for at least 2 months following the last dose of Columvi.

### 2.5.2 Pregnancy

Columvi is not recommended during pregnancy and in women of childbearing potential not using contraception. Female patients of reproductive potential must be advised to avoid pregnancy while receiving Columvi. There are no available data on the use of Columvi in pregnant women. Glofitamab is an immunoglobulin G (IgG). IgG is known to cross the placenta. Based on its mechanism of action, glofitamab is likely to cause fetal B-cell depletion when administered to a pregnant woman. Female patients receiving Columvi should be advised of the potential harm to the fetus. Female patients should be advised to contact the treating physician, should pregnancy occur.

#### Labor and Delivery

The safe use of Columvi during labor and delivery has not been established.

### 2.5.3 Lactation

It is not known whether glofitamab is excreted in human milk. No studies have been conducted to assess the impact of glofitamab on milk production or its presence in human milk. Human IgG is known to be present in human milk. The potential for absorption of glofitamab and the potential for adverse reactions in the nursing infant is unknown. Women should be advised to discontinue breastfeeding during treatment with Columvi and for 2 months after the last dose of Columvi.

## 2.5.4 Pediatric Use

The safety and efficacy of Columvi in pediatric patients have not been established.

## 2.5.5 Geriatric Use

Of the 154 patients with relapsed or refractory DLBCL who were evaluable for safety, 55% were 65 years of age or older. There was a higher rate of fatal adverse events, primarily from COVID-19, in patients 65 years of age or older compared to younger patients. No differences in efficacy of Columvi were observed between patients  $\geq 65$  years of age and those under 65 years. No dose adjustment of Columvi is required in patients  $\geq 65$  years of age. See Section 2.2.1 *Special Dosage Instructions* and Section 3.2.5 *Pharmacokinetics in Special Populations*.

## 2.5.6 Renal Impairment

No dose adjustment is required in patients with mild or moderate renal impairment (CrCL 30 to  $< 90$  mL/min) based on population pharmacokinetic analysis. The safety and efficacy of Columvi in patients with severe renal impairment has not been studied. See Section 2.2.1 *Special Dosage Instructions* and Section 3.2.5 *Pharmacokinetics in Special Populations*.

## 2.5.7 Hepatic Impairment

No dose adjustment is required in patients with mild hepatic impairment (total bilirubin  $>$  ULN to  $\leq 1.5 \times$  ULN or AST  $>$  ULN) based on population pharmacokinetic analysis. The safety and efficacy of Columvi in patients with moderate or severe hepatic impairment has not been studied. See Section 2.2.1 *Special Dosage Instructions* and Section 3.2.5 *Pharmacokinetics in Special Populations*.

## 2.6 UNDESIRABLE EFFECTS

### 2.6.1 Clinical Trials

#### Summary of the safety profile

#### Columvi monotherapy

Approximately 1530 patients with relapsed or refractory non-Hodgkin's lymphoma (NHL) have received Columvi in the clinical development program of Columvi.

The adverse drug reactions described below were identified from clinical studies in patients with relapsed or refractory DLBCL who were treated with Columvi at the recommended dose as monotherapy (N=145) or in combination with gemcitabine and oxaliplatin (N=172).

#### Tabulated summary of adverse drug reactions from clinical trials

Adverse drug reactions from clinical trials (Table 6, Table 7) are listed by MedDRA system organ class. The corresponding frequency category for each adverse drug reaction is based on the following convention: very common ( $\geq 1/10$ ), common ( $\geq 1/100$  to  $< 1/10$ ), uncommon ( $\geq 1/1,000$  to  $< 1/100$ ), rare ( $\geq 1/10,000$  to  $< 1/1,000$ ), very rare ( $< 1/10,000$ ).

**Table 6 Adverse Drug Reactions Occurring in Patients with Relapsed or Refractory DLBCL treated with Columvi Monotherapy**

System Organ Class Adverse Reaction	Columvi N=145		
	All Grades (frequency category)	All Grades (%)	Grade 3-4 (%)
<b>Immune system disorders</b>			
Cytokine release syndrome <sup>a</sup>	Very common	67.6	4.1
<b>Blood and lymphatic system disorders</b>			
Neutropenia <sup>b</sup>	Very common	40.0	29.0
Anaemia <sup>c</sup>	Very common	30.3	7.6
Thrombocytopenia <sup>d</sup>	Very common	24.1	6.9
Lymphopenia <sup>e</sup>	Common	4.8	4.8
Febrile neutropenia <sup>f</sup>	Common	3.4	3.4

General disorders and administration site conditions			
Pyrexia	Very common	15.9	0
<b>Metabolism and nutrition disorders</b>			
Hypophosphataemia	Very common	18.6	6.2
Hypomagnesaemia	Very common	15.2	0
Hypocalcaemia	Very common	12.4	0
Hypokalaemia	Very common	10.3	0.7
Hyponatraemia	Common	8.3	1.4
Tumour lysis syndrome	Common	1.4	1.4
<b>Skin and subcutaneous tissue disorders</b>			
Rash <sup>g</sup>	Very common	20.0	1.4
<b>Gastrointestinal disorders</b>			
Constipation	Very common	14.5	0
Diarrhoea	Very common	13.8	0
Nausea	Very common	10.3	0
Gastrointestinal haemorrhage <sup>h</sup>	Common	2.8	2.8
Vomiting	Common	4.1	0
Colitis	Uncommon	0.7	0.7
<b>Neoplasms benign, malignant and unspecified (incl cysts and polyps)</b>			
Tumour flare	Very common	11.7	2.8
<b>Nervous system disorders</b>			
Headache	Very common	10.3	0
Immune effector cell-associated neurotoxicity syndrome <sup>e,g</sup>	Common	4.8	0.7*
Somnolence	Common	1.4	0.7
Tremor	Common	1.43	0
Myelitis <sup>i</sup>	Uncommon	0.7	0.7
<b>Infections and infestations</b>			
Viral infections <sup>j</sup>	Very common	11.0	3.4*
Bacterial infections <sup>k</sup>	Common	6.2	1.4
Upper respiratory tract infections <sup>l</sup>	Common	5.5	0
Sepsis <sup>m</sup>	Common	4.1	2.8*
Lower respiratory tract infections <sup>n</sup>	Common	2.1	0
Pneumonia	Common	4.1	0.7
Urinary tract infection <sup>o</sup>	Common	2.8	0.7
Fungal infections <sup>p</sup>	Uncommon	1.4	0
<b>Investigations</b>			
Alanine aminotransferase increased	Common	9.0	2.8
Aspartate aminotransferase increased	Common	8.3	2.8
Blood alkaline phosphatase increased	Common	9.0	1.4
Gamma-glutamyltransferase increased	Common	6.9	2.8
Blood bilirubin increased	Common	4.1	0.7
Hepatic enzyme increased	Common	1.4	1.4
<b>Psychiatric disorders</b>			
Confusional state	Common	1.4	0

\* Grade 5 reactions reported include COVID-19 (2.1%), sepsis (1.4%), COVID-19 pneumonia (1.4%), and delirium (0.7%).

a. Based on ASTCT consensus grading.

b. Includes neutropenia and neutrophil count decreased.

c. Includes anaemia and haemoglobin decreased.

d. Includes thrombocytopenia and platelet count decreased.

e. Includes lymphopenia and lymphocyte count decreased.

f. Includes febrile neutropenia and neutropenic infection.

g. Includes rash, rash pruritic, rash maculo-papular, dermatitis, dermatitis acneiform, dermatitis exfoliative, erythema, palmar erythema, pruritus, and rash erythematous.

h. Includes gastrointestinal haemorrhage, large intestinal haemorrhage, and gastric haemorrhage.

i. Myelitis occurred concurrently with CRS.

j. Includes COVID-19, COVID-19 pneumonia, herpes zoster, influenza, and ophthalmic herpes zoster.

k. Includes vascular device infection, bacterial infection, Campylobacter infection, biliary tract infection bacterial, urinary tract infection bacterial, *Clostridium difficile* infection, Escherichia infection, and peritonitis.

l. Includes upper respiratory tract infection, sinusitis, nasopharyngitis, chronic sinusitis, and rhinitis.

m. Includes sepsis and septic shock.

n. Includes lower respiratory tract infection and bronchitis.

o. Includes urinary tract infection and Escherichia urinary tract infection.

p. Includes oesophageal candidiasis and oral candidiasis.

q. Includes somnolence, cognitive disorder, confusional state, delirium, and disorientation.

The most common serious adverse reactions reported in  $\geq 2\%$  of patients were cytokine release syndrome (22.1%), sepsis (3.9%), COVID-19 (3.2%), COVID-19 pneumonia (3.2%) and tumour flare (3.2%).

**Table 7 Adverse Drug Reactions Occurring in Patients with Relapsed or Refractory DLBCL Treated with Columvi in Combination with Gemcitabine and Oxaliplatin**

System Organ Class Adverse Drug Reaction	Columvi+GemOx N=172		
	All Grades (frequency category)	Any Grade (%)	Grade 3-4 (%)
<b>Immune system disorders</b>			
Cytokine release syndrome <sup>a</sup>	Very common	44.2	2.3
<b>Blood and lymphatic system disorders</b>			
Thrombocytopenia <sup>b</sup>	Very common	50.0	26.7
Neutropenia <sup>c</sup>	Very common	44.2	35.5
Anaemia	Very common	41.3	16.9
Lymphopenia <sup>d</sup>	Very common	24.4	12.2
Febrile neutropenia	Common	2.9	2.9
<b>Gastrointestinal Disorders</b>			
Nausea	Very common	41.3	0.6
Diarrhoea	Very common	34.9	3.5
Vomiting	Very common	23.8	0.6
Abdominal pain <sup>e</sup>	Very common	18.6	2.3
Constipation	Very common	18.6	0
Colitis <sup>f</sup>	Common	2.3	1.2
Pancreatitis <sup>g</sup>	Common	1.7	1.2
<b>Nervous system disorders</b>			
Peripheral neuropathy <sup>h</sup>	Very common	37.2	1.2
Headache	Common	8.7	0
ICANS <sup>i</sup>	Common	2.3	0.6
Tremor	Uncommon	0.6	0
<b>Investigations</b>			
Aspartate aminotransferase increased	Very common	34.3	3.5
Alanine aminotransferase increased	Very common	32.6	2.9
Blood alkaline phosphatase increased	Very common	18.0	0.6

Gamma-glutamyltransferase increased <sup>d</sup>	Very common	15.1	2.3
Blood lactate dehydrogenase increased	Very common	11.6	0
Blood bilirubin increased <sup>d</sup>	Common	3.5	0
Hepatic enzyme increased	Uncommon	0.6	0
<b>Skin and subcutaneous disorders</b>			
Rash <sup>h</sup>	Very common	26.7	0.6
<b>General disorders and administration site conditions</b>			
Pyrexia	Very common	24.4	0.6
<b>Musculoskeletal and connective tissue disorders</b>			
Musculoskeletal pain <sup>m</sup>	Very common	18.6	2.3
<b>Metabolism and nutrition disorders</b>			
Hypokalaemia	Very common	18.6	2.9
Hyponatremia	Very common	11.0	0.6
Hypomagnesaemia	Common	7.6	0
Hypocalcaemia	Common	6.4	0.6
Hypophosphataemia	Common	5.8	1.7
Tumour lysis syndrome	Common	1.7	1.7
<b>Infections and infestations</b>			
COVID-19* <sup>n</sup>	Very common	17.4	3.5
Respiratory tract infections* <sup>o</sup>	Very common	13.4	1.7
Pneumonia* <sup>p</sup>	Very common	12.8	5.2
Cytomegalovirus infections <sup>q</sup>	Common	6.4	0.6
Herpes viral infections <sup>r</sup>	Common	5.8	0.6
Urinary tract infection <sup>s</sup>	Common	3.5	1.7
Sepsis* <sup>t</sup>	Common	2.9	2.3
Candida infections <sup>u</sup>	Common	1.7	0
Pneumocystis jirovecii pneumonia	Uncommon	0.6	0.6
<b>Neoplasms benign, malignant and unspecified (incl cysts and polyps)</b>			
Tumour flare <sup>v</sup>	Common	2.3	0
<b>Respiratory, thoracic, and mediastinal disorders</b>			
Pneumonitis* <sup>w</sup>	Common	1.2	0

\* Grade 5 reactions reported include COVID-19 (1.7%), pneumonitis (1.2%), septic shock (0.6%), pneumonia (0.6%), and respiratory tract infection (0.6%).

- Based on ASTCT consensus grading.
- Includes thrombocytopenia and platelet count decreased.
- Includes neutropenia and neutrophil count decreased.
- Includes lymphopenia and lymphocyte count decreased.
- Includes abdominal pain, abdominal discomfort, abdominal pain upper, abdominal pain lower, and gastrointestinal pain.
- Includes colitis, colitis ischaemic, and enterocolitis.
- Includes pancreatitis and pancreatitis acute.
- Includes neuropathy peripheral, peripheral sensory neuropathy, dysaesthesia, paraesthesia, hypoesthesia, peripheral motor neuropathy, and polyneuropathy.
- Includes confusional state, delirium, and ICANS.
- Includes gamma-glutamyltransferase increased and gamma-glutamyltransferase abnormal.
- Includes blood bilirubin increased and hyperbilirubinaemia.
- Includes rash, rash pruritic, rash maculo-papular, erythema, pruritus, rash erythematous, urticaria, and erythema multiforme.
- Includes arthralgia, musculoskeletal pain, back pain, bone pain, myalgia, neck pain, pain in extremity, musculoskeletal chest pain, and non-cardiac chest pain.
- Includes COVID-19, COVID-19 pneumonia, and SARS-CoV-2 test positive.
- Includes upper respiratory tract infection, lower respiratory tract infection, respiratory tract infection, and respiratory tract infection bacterial.
- Includes pneumonia, pneumonia bacterial, and pneumonia pneumococcal.
- New onset or reactivation. Includes cytomegalovirus infection, cytomegalovirus test positive, cytomegalovirus infection reactivation, and cytomegalovirus viraemia.
- New onset or reactivation. Includes herpes zoster and herpes virus infection.
- Includes urinary tract infection and urosepsis.
- Includes sepsis, streptococcal sepsis, septic shock, and enterococcal sepsis.
- Includes oral candidiasis and candida infection.
- Includes tumour flare and tumour pain.

#### Description of selected adverse drug reactions from clinical trials

The descriptions below reflect information for significant adverse reactions for Columvi monotherapy and/or combination therapy. Details for the significant adverse reactions for Columvi when given in combination are presented separately if clinically relevant differences were noted in comparison to Columvi monotherapy.

#### Cytokine Release Syndrome

##### Columvi monotherapy

Any grade CRS (by ASTCT criteria) occurred in 67.6% of patients who received Columvi monotherapy, with Grade 1 CRS reported in 50.3% of patients, Grade 2 CRS in 13.1% of patients, Grade 3 CRS in 2.8% of patients, and Grade 4 CRS in 1.4% of patients. CRS occurred more than once in 32.4% (47/145) of patients; 36/47 patients experienced multiple Grade 1 CRS events only.

There were no fatal cases of CRS. CRS resolved in all patients except one. One patient discontinued Columvi due to CRS.

In patients with CRS, the most common manifestations of CRS included pyrexia (99.0%), tachycardia (25.5%), hypotension (23.5%), chills (14.3%), and hypoxia (12.2%). Grade 3 or higher events associated with CRS included hypotension (3.1%), hypoxia (3.1%), pyrexia (2.0%), and tachycardia (2.0%).

CRS of any grade occurred in 54.5% of patients following the 2.5 mg dose of Columvi at Cycle 1 Day 8 with median time to onset (from the start of infusion) of 12.6 hours (range: 5.2 to 50.8 hours) and median duration of 31.8 hours (range: 0.5 to 316.7 hours); in 33.3% of patients following the 10 mg dose at Cycle 1 Day 15 with median time to onset of 26.8 hours (range: 6.7 to 125.0 hours) and median duration of 16.5 hours (range: 0.3 to 109.2 hours); and in 26.8% of patients following the 30 mg dose at Cycle 2 Day 1 with median time to onset of 28.2 hours (range: 15.0 to 44.2 hours) and median duration of 18.9 hours (range: 1.0 to 180.5 hours). CRS was reported in 0.9% of patients at Cycle 3 and in 2% of patients beyond Cycle 3.

In 145 patients, 7 (4.8%) patients experienced elevated liver function tests (AST and ALT > 3 ULN and/or total bilirubin > 2 ULN) reported concurrently with CRS (n=6) or with disease progression (n=1).

Hospitalisations due to patients experiencing CRS following Columvi administration occurred in 22.1% of patients and the reported median duration of hospitalisation was 4 days (range: 2 to 15 days).

Grade ≥ 2 CRS occurred in 12.4% of patients following the first Columvi dose (2.5 mg), with median time to onset of 9.7 hours (range: 5.2 to 19.1 hours) and median duration of 50.4 hours (range: 6.5 to 316.7 hours). Following Columvi 10 mg dose at Cycle 1 Day 15, the incidence of Grade ≥ 2 CRS decreased to 5.2% of patients, with median time to onset of 26.2 hours (range: 6.7 to 144.2 hours) and median duration of 30.9 hours (range: 3.7 to 227.2 hours). Grade ≥ 2 CRS following Columvi 30 mg dose at Cycle 2 Day 1 occurred in one patient (0.8%) with time to onset of 15.0 hours and duration of 44.8 hours. No Grade ≥ 2 CRS was reported beyond Cycle 2.

Among the 25 patients who experienced Grade 2 or higher CRS after Columvi, 22 (88%) received tocilizumab, 15 (60%) received corticosteroids, and 14 (56%) received both tocilizumab and corticosteroids. Ten patients (40%) received oxygen. All 6 patients (24.0%) with Grade 3-4 CRS received a single vasopressor.

In patients who received dexamethasone premedication (N=39) versus another glucocorticoid premedication (n=106), CRS of any grade occurred in 48.7% vs. 56.6% of patients; Grade 1 CRS in 38.5% vs. 43.4% of patients; Grade 2 CRS in 7.7% vs. 9.4% of patients; Grade 3 CRS in 2.6% vs. 1.9% of patients; and Grade 4 CRS in 0% vs. 1.9% of patients after the 2.5 mg dose of Columvi at Cycle 1 Day 8. After the 10 mg dose at Cycle 1 Day 15 (n=36 for dexamethasone premedication, n=99 for another glucocorticoid premedication), any grade CRS occurred in 22.2% vs 37.4% of patients; Grade 1 CRS in 22.2% vs 30.3% of patients; Grade 2 CRS in 0% vs 6.1% of patients; and Grade 3 CRS in 0% vs 1% of patients. After the 30 mg dose at Cycle 2 Day 1 (n=32 for dexamethasone premedication, n=95 for another glucocorticoid premedication) any grade CRS occurred in 6.3% vs 33.7% of patients; Grade 1 CRS in 6.3% vs 32.6% of patients; and Grade 2 CRS in 0% vs 1.1% of patients.

#### Columvi in combination with gemcitabine and oxaliplatin

Any grade CRS (by ASTCT criteria) occurred in 44.2% of patients who received Columvi with gemcitabine and oxaliplatin, with Grade 1 CRS reported in 31.4% of patients, Grade 2 CRS in 10.5% of patients, and Grade 3 CRS in 2.3% of patients. CRS occurred more than once in 21.5% (37/172) of patients; 30/37 patients experienced multiple Grade 1 CRS events only. There were no Grade 4 or fatal cases of CRS. CRS resolved in all patients except one. One patient discontinued Columvi due to CRS.

In patients with CRS, the most common manifestations of CRS included pyrexia (98.7%), hypotension (22.4%), chills (17.1%) and hypoxia (14.5%). Grade 3 or higher events associated with CRS included hypotension (6.6%), hypoxia (5.3%), pyrexia (3.9%), chills (1.3%), and diarrhea (1.3%).

CRS of any grade occurred in 34.9% of patients following the 2.5 mg dose of Columvi at Cycle 1 Day 8 with median time to onset (from the start of infusion) of 12.6 hours (range: 4.4 to 54.7 hours) and median duration of 19.8 hours (range: 2.0 to 168.0 hours); in 14.4% of patients following the 10 mg dose at Cycle 1 Day 15 with median time to onset of 22.8 hours (range: 7.4 to 81.2 hours) and median duration of 10.6 hours (range: 1.0 to 248.5 hours); and in 9.3% of patients following the 30 mg dose at Cycle 2 Day 1 with median time to onset of 23.5 hours (range: 14.7 to 33.4 hours) and median duration of 18.4 hours (range: 8.3 to 137.0 hours). CRS was reported in 6.7% of patients at Cycle 3 and in 11.0% of patients beyond Cycle 3.

Grade ≥ 2 CRS occurred in 10.5% of patients following the first Columvi dose (2.5 mg), with median time to onset of 12.0 hours (range: 4.4 to 30.5 hours) and median duration of 42.3 hours (range: 3.5 to 143.7 hours). The majority (14/18) of patients who experienced Grade ≥ 2 CRS had onset of CRS within 8 hours of the start of the first Columvi dose (2.5 mg) or presented with a fever ≥ 1.5 hours before the onset of other symptoms of a Grade ≥ 2 CRS. Following Columvi 10 mg dose at Cycle 1 Day 15, the incidence of Grade ≥ 2 CRS decreased to 1.8% of patients, with median time to onset of 22.3 hours (range: 7.4 to 22.8 hours) and median duration of 37.0 hours (range: 34.8 to 248.5 hours). There were no Grade ≥ 2 CRS events following Columvi 30 mg dose at Cycle 2 Day 1. Three patients (2%) had Grade ≥ 2 CRS beyond Cycle 2 (all Grade 2 events).

Of the 172 patients, 2 patients (1.2%) experienced elevated liver function tests (AST and ALT > 3 × ULN) reported concurrently with CRS.

Out of the 76 patients with any grade CRS, 28 patients (36.8%) were treated with tocilizumab, 39 patients (51.3%) were treated with corticosteroids, and 18 patients (23.7%) received both tocilizumab and corticosteroids.

Among the 22 patients who experienced Grade 2 or higher CRS after Columvi, 16 (72.7%) received tocilizumab, 15 (68.2%) received corticosteroids, and 12 (54.5%) received both tocilizumab and corticosteroids. Eleven patients (50.0%) received oxygen. All 4 patients (18.2%) with Grade 3 CRS received a single vasopressor.

Hospitalisations due to patients experiencing CRS following Columvi administration occurred in 19.8% of patients and the reported median duration of hospitalisation was 5 days (range: 2 to 85 days).

#### Neurologic Toxicity

##### Columvi monotherapy

The most frequent neurologic toxicities of any grade were headache (9.7%), peripheral neuropathy (1.9%), dizziness or vertigo (6.5%), and mental status changes (5.8%), including confusional state, cognitive disorder, disorientation, somnolence, and delirium). Grade 3 or higher neurologic adverse reactions occurred in 2.6% of patients and included somnolence, delirium, and myelitis. Cases of ICANS of any grade occurred in 4.8% of patients treated with Columvi.

##### Columvi in combination with gemcitabine and oxaliplatin

Neurologic toxicity was reported in 59.3% of patients who received Columvi with gemcitabine and oxaliplatin. The most frequent neurologic toxicities of any grade were peripheral neuropathy (37.2%), insomnia (11.6%), headache (8.7%), dizziness (7.0%), and dysgeusia (5.2%). Grade 3 or higher neurologic adverse reactions occurred in 5.8% of patients and included peripheral sensory neuropathy, syncope, extrapyramidal disorder, confusional state, delirium, herpes zoster, meningitis aseptic, subdural hematoma, and cerebral hemorrhage.

Four patients (2.3%) experienced ICANS: 1 patient (0.6%) experienced a Grade 1 event (confusional state), 2 patients (1.2%) experienced a Grade 2 event (1 patient with confusional state, and 1 patient with ICANS), and 1 patient (0.6%) experienced Grade 3 delirium.

The onset of the 4 events of ICANS was concurrent with CRS. The median time to resolution of ICANS was 1.5 days (range: 1 to 4 days).

#### Serious Infections

Serious infections were reported in 15.9% of patients who received Columvi monotherapy. The most frequent serious infections reported in ≥ 2% of patients were sepsis (4.1%), COVID-19 (3.4%), and COVID-19 pneumonia (2.8%). Infection-related deaths were reported in 4.8% of patients (due to sepsis, COVID-19 pneumonia, and COVID-19). Four patients (2.8%) experienced serious infections concurrently with Grade 3-4 neutropenia.

Serious infections were reported in 22.7% of patients who received Columvi with gemcitabine and oxaliplatin. The most frequent serious infections reported in ≥ 2% of patients were pneumonia (5.8%), COVID-19 (4.7%), and lower respiratory tract infection (2.9%). Infection-related deaths were reported in 3.5% of patients (due to COVID-19, pneumonia, respiratory tract infection, and septic shock). One patient (0.6%) experienced a serious infection (pneumonia) concurrently with Grade 3 neutropenia.

#### Pneumonitis

Pneumonitis events (excluding pneumonia of infectious etiology) were reported in 2 patients (1.2%) who received Columvi with gemcitabine and oxaliplatin, both of which were fatal events. The median time to onset of pneumonitis from the first Columvi dose was 168 days (range: 102 to 255 days).

#### Colitis

Colitis (Grade 4) was reported in 1 patient who received Columvi monotherapy, with time to onset from the first Columvi dose of 104 days.

Colitis events (excluding infectious etiology) were reported in 4 patients (2.3%) who received Columvi with gemcitabine and oxaliplatin. Two patients (1.2%) had Grade 3 events. The median time to onset of colitis from the first Columvi dose was 154 days (range: 115 to 187 days).

#### Opportunistic Infections

CMV events were reported in 6/467 patients (1.3%) who received Columvi monotherapy in study NP30179, with 1 patient (0.2%) experiencing Grade 3 CMV chorioretinitis. Pneumocystis jirovecii pneumonia was reported in 4/467 patients (0.9%), 3 of whom (0.6%) had Grade 3 events.

CMV events were reported in 11 patients (6.4%) who received Columvi with gemcitabine and oxaliplatin, with 1 patient (0.6%) experiencing Grade 3 CMV viraemia. Oral candidiasis was reported in 3 patients (1.7%), all of which were Grade 1-2 events. Pneumocystis jirovecii pneumonia (Grade 3) was reported in 1 patient (0.6%), the same patient with Grade 3 CMV viraemia. Borellia meningitis (Grade 2) was reported in 1 patient (0.6%).

#### Neutropenia

Neutropenia (including neutrophil count decreased) was reported in 40.0% of patients and severe neutropenia (Grade 3-4) was reported in 29.0% of patients who received Columvi monotherapy. The median time to onset of the first neutropenia event was 29 days (range: 1 to 203 days). Prolonged neutropenia (lasting longer than 30 days) occurred in 11.7% of patients. The majority of patients with neutropenia (79.3%) were treated with G-CSF. Febrile neutropenia was reported in 3.4% of patients.

#### Tumor Flare

Tumor flare was reported in 11.7% of patients who received Columvi monotherapy, including Grade 2 tumor flare in 4.8% of patients and Grade 3 tumor flare in 2.8% of patients. Tumor flare was reported involving lymph nodes in the head and neck presenting with pain, and involving lymph nodes in the thorax with symptoms of breathlessness due to development of pleural effusion. Most tumor flare events (16/17) occurred during Cycle 1, and no tumor flare events were reported beyond Cycle 2. The median time to onset of tumor flare of any grade was 2 days (range: 1 to 16 days), and

the median duration was 3.5 days (range: 1 to 35 days). Among the 11 patients who experienced Grade ≥ 2 tumour flare, of which 2 (18.2%) patients received analgesics, 6 (54.5%) patients received corticosteroids and analgesics including morphine derivatives, 1 (0.9%) patient received corticosteroids and anti-emetics, and 2 (18.2%) patients did not require treatment. All tumour flare events resolved except in one patient\* with a Grade ≥ 2 event. No patients discontinued Columvi due to tumor flare.

\*One patient had tumour flare unresolved at the time of death due to disease progression. The treatment of tumour flare for this patients was not reported.

#### Tumor Lysis Syndrome

TLS was reported in 2 patients (1.4%) who received Columvi monotherapy and was Grade 3 in severity in both cases. The median time to onset of TLS was 2 days, and the median duration was 4 days (range: 3 to 5 days).

#### Laboratory Abnormalities

Table 8 summarizes treatment-emergent shifts from baseline in laboratory abnormalities with Columvi monotherapy. Table 9 summarizes treatment-emergent shifts from baseline in laboratory abnormalities with Columvi in combination with gemcitabine and oxaliplatin.

**Table 8 Laboratory Abnormalities Worsening from Baseline, with Grade 3 to 4 Occurring in ≥ 10% of Patients with Relapsed or Refractory DLBCL Treated with Columvi Monotherapy**

Laboratory Abnormality <sup>a</sup>	Columvi NCI CTCAE Grade	
	All Grades (%) <sup>b</sup>	Grade 3 or 4 (%) <sup>b,c</sup>
<b>Hematology</b>		
Decreased lymphocytes	90.2	83.2
Decreased neutrophils	55.6	25.7
Decreased leukocytes	71.0	13.8
<b>Chemistry</b>		
Hypophosphatemia	68.8	27.8
Hyperglycemia	14.2	14.2
Hyperuricemia	22.6	22.6

a. Percentages based on patients with a baseline and at least one post-baseline assessment for the specific laboratory parameter.

b. N=143 for decreased lymphocytes; N=144 for decreased neutrophils; N=145 for decreased leukocytes; N=144 for hypophosphatemia; N=141 for hyperglycemia; N=137 for hyperuricemia.

c. Includes shifts from NCI CTCAE Grade 0-2 at baseline to Grade ≥ 3 post-baseline, and shifts from Grade 3 at baseline to Grade 4 post-baseline.

**Table 9 Laboratory Abnormalities Worsening from Baseline, with Grade 3 to 4 Occurring in ≥ 5% of Patients with Relapsed or Refractory DLBCL Treated with Columvi in Combination with Gemcitabine and Oxaliplatin**

Laboratory Abnormality <sup>a</sup>	Columvi+GemOx NCI CTCAE Grade	
	All Grades (%) <sup>b</sup>	Grade 3 or 4 (%) <sup>b,c</sup>
<b>Hematology</b>		
Decreased platelets	82.6	29.7
Decreased lymphocytes	81.1	50.3
Decreased hemoglobin	73.8	19.8
Decreased neutrophils	65.3	39.5
Decreased leukocytes	58.1	26.2
<b>Chemistry</b>		
Hyperuricemia	21.9	21.9
Hypokalemia	23.3	5.8

a. Percentages based on patients with a baseline and at least one post-baseline assessment for the specific laboratory parameter.

b. N=172 for decreased platelets; N=143 for decreased lymphocytes; N=172 for decreased hemoglobin; N=147 for decreased neutrophils; N=172 for decreased leukocytes; N=160 for hyperuricemia; N=172 for hypokalemia.

c. Includes shifts from Grade 0-2 at baseline to Grade ≥ 3 post-baseline, and shifts from Grade 3 at baseline to Grade 4 post-baseline.

#### 2.6.2 Post marketing Experience

Not applicable.

#### 2.7 OVERDOSE

There is no experience with overdose of Columvi in clinical trials. In case of overdose, patients should be closely monitored for signs and symptoms of adverse reactions, and appropriate symptomatic treatment instituted.

#### 2.8 INTERACTIONS WITH OTHER MEDICINAL PRODUCTS AND OTHER FORMS OF INTERACTION

No clinical drug-drug interaction studies have been performed.

No drug interactions with Columvi are expected via the cytochrome P450 enzymes, other metabolizing enzymes, or transporters.

For certain CYP substrates (e.g. warfarin, cyclosporine) where minimal concentration changes may lead to serious adverse reactions, monitor for toxicities or drug concentrations of such CYP substrates when coadministered with Columvi.

Glofitamab causes the release of cytokines that may suppress the activity of CYP enzymes, resulting in increased exposure of CYP substrates. Increased exposure of CYP substrates is more likely to occur after the first dose of Columvi on Cycle 1 Day 8 and up to 14 days after the first 30 mg dose on Cycle 2 Day 1 and during and after CRS.

The pharmacokinetics of glofitamab are not affected by co-administration with gemcitabine or oxaliplatin.

#### 3. PHARMACOLOGICAL PROPERTIES AND EFFECTS

##### 3.1 PHARMACODYNAMIC PROPERTIES

###### 3.1.1 Mechanism of Action

Glofitamab is a bispecific monoclonal antibody that binds bivalently (with high avidity) to CD20 expressed on the surface of B cells, and monovalently to CD3 in the T-cell receptor complex expressed on the surface of T cells. By simultaneous binding to CD20 on the B cell and CD3 on the T cell, glofitamab mediates the formation of an immunological synapse with subsequent potent T-cell activation and proliferation, secretion of cytokines, and release of cytolytic proteins that results in the lysis of CD20-expressing B cells.

##### Pharmacodynamics

In study NP30179, peripheral B-cell counts were < 70 cells/μL in almost all patients (98.6%) with relapsed and refractory LBCL prior to Columvi treatment initiation and remained low during Columvi treatment.

During Cycle 1 (step-up dosing), transient increases in plasma IL-6 levels were observed at 6 hours post-Columvi infusion, which remained elevated at 20 hours post-infusion and returned to baseline prior to the next infusion.

In study GO41944 (STARGLO), peripheral B-cell counts were <70 cells/μL in the majority of patients (79.4%) with relapsed and refractory DLBCL prior to Columvi treatment initiation and remained low during Columvi treatment.

##### Cardiac electrophysiology

In study NP30179, 16/145 patients who were exposed to glofitamab experienced a post-baseline QTc value > 450ms. One of these cases was assessed to be of clinical significance by the investigator. No patients discontinued treatment due to QTc prolongation.

##### 3.1.2 Clinical / Efficacy Studies Relapsed or Refractory DLBCL

###### Columvi monotherapy

The efficacy of Columvi monotherapy was evaluated in study NP30179, a single-arm, open-label, multicenter, multi-cohort trial, which included 155 patients with relapsed or refractory DLBCL after at least two prior lines of systemic therapy including an anti-CD20 monoclonal antibody and an anthracycline agent. The study excluded patients with prior allogeneic hematopoietic stem cell transplant, previous or active central

nervous system (CNS) lymphoma, active infection, recent infection requiring intravenous antibiotics, ECOG performance status  $\geq 2$ , creatinine clearance (CrCL)  $< 50$  mL/min, or hepatic transaminases  $> 3 \times$  ULN.

Following pre-treatment with obinutuzumab at Cycle 1 Day 1, patients received 2.5 mg of Columvi at Cycle 1 Day 8, 10 mg of Columvi at Cycle 1 Day 15, and 30 mg of Columvi at Cycle 2 Day 1 as per the step-up dosing schedule. Patients continued to receive 30 mg of Columvi on Day 1 of Cycles 3 to 12. Patients received premedication including an anti-pyretic, an anti-histamine and a glucocorticoid (see section 2.2 *Dosage and Administration*). The duration of each cycle was 21 days.

Patients received a median of 5 cycles of Columvi treatment (range: 1 to 13 cycles).

The baseline demographic and disease characteristics were: median age 66 years (range: 21 to 90 years); 65.2% males; 76.8% white, 4.5% Asian, and 1.9% Black or African American; 5.8% Hispanic or Latino; and ECOG performance status of 0 (44.5%) or 1 (54.2%). Most patients (71.0%) had DLBCL not otherwise specified, 18.7% had DLBCL transformed from follicular lymphoma, 6.5% had high-grade B-cell lymphoma (HGBCL), and 3.9% had PMBCL. The median number of prior lines of therapy was 3 (range: 2 to 7); 39.4% of patients received 2 prior lines and 60.6% received 3 or more prior lines of therapy. All patients had received prior chemotherapy and anti-CD20 monoclonal antibody therapy; 33.5% of patients had received prior CAR T-cell therapy, and 18.1% of patients had received autologous stem cell transplant. Most patients (89.7%) had refractory disease, 58.7% patients had primary refractory disease, 84.5% of patients were refractory to their last prior therapy, and 88.5% of patients who received prior CAR T-cell therapy were refractory to CAR T-cell therapy.

The overall median duration of follow-up was 13.4 months (range: 0 to 28 months). Median duration of follow-up from the date of first response per Independent Review Committee (IRC) assessment was 12.0 months (range: 0 to 27 months).

The primary efficacy outcome measure was complete response (CR) rate as assessed by IRC using 2014 Lugano criteria. The secondary efficacy outcome measures included Investigator (INV)-assessed CR, and overall response rate (ORR), duration of response (DOR), duration of complete response (DOCR), time to first response (TFOR), time to first complete response (TFCR), overall survival (OS), and progression-free survival (PFS), as assessed by IRC and by INV.

Efficacy results are summarized in Table 10.

**Table 10 Efficacy in Patients with Relapsed or Refractory DLBCL Treated with Columvi Monotherapy**


Efficacy Endpoints	Columvi N=155	
<b>Primary Endpoint</b>		
<b>IRC-Assessed Complete Response</b>		
Patients with CR, n (%)	62 (40.0)	
95% CI	[32.22, 48.17]	
<b>Secondary Endpoints</b>		
<b>INV-Assessed Complete Response</b>		
Patients with CR, n (%)	59 (38.1)	
95% CI	[30.39, 46.20]	
<b>Overall Response Rate</b>	<i>IRC-Assessed</i>	<i>INV-Assessed</i>
Patients with CR or PR, n (%)	80 (51.6)	91 (58.7)
95% CI	[43.46, 59.70]	[50.53, 66.55]
Partial Response (PR), n (%)	18 (11.6)	32 (20.6)
95% CI	[7.03, 17.73]	[14.57, 27.88]
<b>Duration of Complete Response<sup>a</sup></b>	<i>IRC-Assessed</i>	<i>INV-Assessed</i>
Median DOCR, months [95% CI]	NE [16.8, NE]	NE [19.8, NE]
Range, months	0 <sup>b</sup> -27 <sup>b</sup>	0 <sup>b</sup> -27 <sup>b</sup>
9-month DOCR, % [95% CI] <sup>c</sup>	76.0 [63.26, 88.71]	72.5 [59.25, 85.68]
12-month DOCR, % [95% CI] <sup>c</sup>	73.1 [59.57, 86.53]	72.5 [59.25, 85.68]
<b>Duration of Response<sup>d</sup></b>	<i>IRC-Assessed</i>	<i>INV-Assessed</i>
Median DOR, months [95% CI]	16.8 [10.4, NE]	10.4 [5.4, NE]
Range, months	0 <sup>b</sup> -27 <sup>b</sup>	0 <sup>b</sup> -27 <sup>b</sup>
9-month DOR, % [95% CI] <sup>c</sup>	66.5 [54.91, 78.00]	52.2 [41.10, 63.34]
12-month DOR, % [95% CI] <sup>c</sup>	59.6 [46.85, 72.28]	48.4 [36.93, 59.91]
<b>Time to First Response</b>	<i>IRC-Assessed</i>	<i>INV-Assessed</i>
Median TFOR, days [95% CI]	42 [41, 42]	42 [40, 42]
Range, days	31-178	31-178
<b>Time to First Complete Response</b>	<i>IRC-Assessed</i>	<i>INV-Assessed</i>
Median TFCR, days [95% CI]	42 [42, 44]	43 [42, 48]
Range, days	31-308	31-274
<b>Progression-Free Survival</b>	<i>IRC-Assessed</i>	<i>INV-Assessed</i>
Patients with event, n (%)	95 (61.3)	98 (63.2)
Median PFS, months [95% CI]	4.9 [3.4, 8.1]	3.8 [3.3, 5.4]
6-month PFS, % [95% CI] <sup>c</sup>	46.7 [38.40, 54.92]	39.1 [30.98, 47.14]
9-month PFS, % [95% CI] <sup>c</sup>	39.6 [31.34, 47.76]	35.1 [27.08, 43.03]
12-month PFS, % [95% CI] <sup>c</sup>	34.9 [26.48, 43.31]	30.6 [22.55, 38.69]
<b>Overall Survival</b>	<i>IRC-Assessed</i>	<i>INV-Assessed</i>
Patients with event, n (%)	81 (52.3)	81 (52.3)
Median OS, months [95% CI]	12 [8.0, 16.1]	12 [8.0, 16.1]
6-month OS, % [95% CI] <sup>c</sup>	71.6 [64.34, 78.89]	71.6 [64.34, 78.89]
9-month OS, % [95% CI] <sup>c</sup>	54.8 [46.65, 62.87]	54.8 [46.65, 62.87]
12-month OS, % [95% CI] <sup>c</sup>	50.4 [42.06, 58.71]	50.4 [42.06, 58.71]

CI=confidence interval; INV=Investigator; IRC=Independent Review Committee; N/A=not applicable; NE=not estimable.


a. From date of first complete response until disease progression or death due to any cause.  
b. Censored observations.  
c. Event-free rates based on Kaplan-Meier estimates.  
d. From date of first response (PR or CR) until disease progression or death due to any cause.

The efficacy population included a cohort of patients (N=40) where dexamethasone was mandated as the glucocorticoid premedication. In this cohort, the IRC-assessed ORR was 52.5% (95% CI: 36.1, 68.5) and the CR rate was 47.5% (95% CI: 31.5, 63.9).

**Figure 1 Duration of IRC-Assessed Complete Response in Patients with Relapsed or Refractory DLBCL Treated with Columvi Monotherapy**



**Figure 2 Duration of IRC-Assessed Response in Patients with Relapsed or Refractory DLBCL Treated with Columvi Monotherapy**



The efficacy results based on the non-Hodgkin lymphoma subtype are summarized in Table 11.

**Table 11 IRC-Assessed Response Rates by Histology Subtypes for Patients with Relapsed or Refractory DLBCL Treated with Columvi Monotherapy**

Efficacy Endpoints	DLBCL NOS n=110	Transformed Follicular Lymphoma n=29	High Grade B-cell Lymphoma n=10	Primary mediastinal B-cell lymphoma n=6
CRR, n(%)	44 (40.0%)	14 (48.3%)	1 (10.0%)	3 (50.0%)
95% CI	[30.8, 49.8]	[29.5, 67.5]	[0.3, 44.5]	[11.8, 88.2]
ORR, n(%)	57 (51.8%)	17 (58.6%)	2 (20.0%)	4 (66.7%)
95% CI	[42.1, 61.5]	[38.9, 76.5]	[2.5, 55.6]	[22.3, 95.7]

**Patient-Reported Outcomes**

Study NP30179 evaluated patient-reported outcomes of Columvi treatment. Patients reported moderate to moderate-high levels of Physical Functioning, Role Functioning, and Global Health Status/QoL and low levels of fatigue (weakness, tiredness) as measured by the EORTC QLQ-C30 at baseline which were maintained during treatment. Most patients indicated that symptoms commonly associated with Columvi treatment (constipation, diarrhea, and nausea) were not present or were of low severity if present, and maintained during treatment. Patients reported low levels of lymphoma symptoms at baseline as measured by the FACT-Lym scale which were maintained during treatment. The results should be interpreted with caution in the context of the open-label study design.

**Columvi in combination with gemcitabine and oxaliplatin**

The efficacy of Columvi in combination with gemcitabine and oxaliplatin (Columvi+GemOx) was evaluated in study GO41944 (STARGLO), an open-label multicenter, randomized clinical trial that included 274 patients with relapsed or refractory DLBCL not otherwise specified (DLBCL NOS). The study excluded patients who received only one prior line of therapy who were candidates for stem cell transplant, patients with HGBCL, PMBCL, history of transformation of indolent disease to DLBCL, prior allogeneic hematopoietic stem cell transplant, previous or active CNS lymphoma, ECOG performance status  $> 2$ , CrCL  $< 30$  mL/min, or hepatic transaminases  $> 2.5 \times$  ULN.

Patients who received only one prior line of therapy were not considered candidates for transplant if they met at least one of the following criteria: age  $\geq 70$  years, ECOG performance status  $\geq 2$ , left ventricular ejection fraction  $\leq 40\%$ , insufficient response to salvage therapy prior to ASCT, CrCL  $\leq 45$  mL/min, other comorbidities or criteria that preclude use of transplant based on local practice standards or in the investigator's opinion, or patient refusal of high-dose chemotherapy and/or transplant. Patients were randomized 2:1 to receive Columvi+GemOx (N=183) or rituximab in combination with gemcitabine and oxaliplatin (R-GemOx; N=91) for 8 cycles, followed by 4 additional cycles of Columvi monotherapy for patients in the Columvi+GemOx arm. Randomization was stratified by number of previous lines of systemic therapy for DLBCL (1 vs.  $\geq 2$ ) and outcome of last systemic therapy (relapsed vs. refractory).

In the Columvi+GemOx arm, following pre-treatment with obinutuzumab at Cycle 1 Day 1, patients received 2.5 mg of Columvi at Cycle 1 Day 8, 10 mg of Columvi at Cycle 1 Day 15, and 30 mg of Columvi at Cycle 2 Day 1 as per the step-up dosing schedule. Patients continued to receive 30 mg of Columvi on Day 1 of Cycles 3 to 12. Patients received premedication including an anti-pyretic, an anti-histamine, and dexamethasone. For patients who were unable to receive dexamethasone, an alternate corticosteroid was used (see section 2.2 *Dosage and Administration*). Gemcitabine (1000 mg/m<sup>2</sup>) and oxaliplatin (100 mg/m<sup>2</sup>) were administered intravenously on Day 2 of Cycle 1 and then on Day 1 of subsequent cycles, up to Cycle 8. The duration of each cycle was 21 days.

Patients received a median of 11 cycles of Columvi treatment (range: 1 to 13 cycles).

The baseline demographic and disease characteristics were: median age 68 years (range: 20 to 88 years); 57.7% males; 42.0% white, 50.0% Asian, and 1.1% Black or African American; 5.8% Hispanic or Latino; and ECOG performance status of 0 (43.3%), 1 (46.6%), or 2 (10.1%). The majority of patients (62.8%) had received 1 prior line of systemic therapy; 37.2% of patients received 2 or more prior lines. All patients had received prior chemotherapy and most (98.5%) had received anti-CD20 monoclonal antibody therapy; 7.7% of patients had received CAR T-cell therapy, and 4.0% had received autologous stem cell transplant. The majority of patients (66.8%) had refractory disease; 55.8% had primary refractory disease, and 60.6% of patients were refractory to their last prior therapy. The most common reasons why patients were not considered candidates for transplant included age (42.3%), patient refused high-dose chemotherapy and/or transplant (34.7%), and insufficient response to salvage therapy (9.9%).

The primary efficacy outcome measure was overall survival (OS). At the time of the prespecified primary analysis, a statistically significant improvement in OS was observed for patients randomized to the Columvi+GemOx arm compared to patients randomized to R-GemOx. Statistically significant improvements in PFS and CR rate, as assessed by an IRC, were also observed with Columvi+GemOx over R-GemOx.

Overall survival, PFS, and CR results from an updated analysis conducted after an additional 10.5 months of follow-up continue to demonstrate benefit of Columvi+GemOx over R-GemOx. Median OS was 25.5 months in the Columvi+GemOx arm compared with 12.9 months in the R-GemOx arm in the updated analysis. The 24-month OS rate was 52.8% (95% CI: 44.8, 60.7) in the Columvi+GemOx arm and 33.5% (95% CI: 22.2, 44.9) in the R-GemOx arm.

Pre-specified exploratory subgroup analyses raised a hypothesis of differential efficacy across geographical regions but were not powered to detect statistical significance.

The observed efficacy results by region should be carefully interpreted taking into consideration confounding factors such as small sample sizes relative to the ITT population, high variability and broad confidence intervals for hazard ratios (HRs), which reflect uncertainty in these estimates (see Table 12).

**Table 12 Summary of Key Results Based on Pre-Specified Regional Subgroups**

	Europe (N=88)		North America (N=25)		Rest Of World (N=161)	
	Glofit-GemOx (N=62)	R-GemOx (N=26)	Glofit-GemOx (N=15)	R-GemOx (N=10)	Glofit-GemOx (N=106)	R-GemOx (N=55)
<b>OS</b>						
Median OS, months (95% CI)	21.2 (10.5, NE)	13.8 (11.1, NE)	13.3 (5.2, NE)	NE (7.5, NE)	NE (20.4, NE)	8.3 (5.5, 14.5)
Unstratified HR (95% CI)	1.09 (0.54, 2.18)		2.62 (0.56, 12.34)		0.41 (0.27, 0.64)	
<b>IRC-assessed PFS</b>						
Median IRC PFS, months (95% CI)	9.2 (6.1, 17.0)	7.8 (2.6, NE)	7.5 (2.5, NE)	27.1 (3.3, NE)	20.5 (9.3, NE)	2.5 (1.5, 5.2)
Unstratified HR (95% CI)	0.84 (0.44, 1.59)		2.25 (0.48, 10.54)		0.27 (0.17, 0.42)	

Efficacy results are summarized in Table 13.


**Table 13 Efficacy in Patients with Relapsed or Refractory DLBCL Treated with Columvi in Combination with Gemcitabine and Oxaliplatin (ITT)**

	Primary Analysis		Updated Analysis	
	Median observation time=11.3 months	Median observation time=20.7 months	Median observation time=11.3 months	Median observation time=20.7 months
	Columvi+GemOx (N=183)	R-GemOx (N=91)	Columvi+GemOx (N=183)	R-GemOx (N=91)
<b>Primary Endpoint</b>				
<b>Overall Survival</b>				
Number (%) of deaths	61 (33.3)	40 (44.0)	80 (43.7)	52 (57.1)


Median [95% CI], months	NE [13.8, NE]	9.0 [7.3, 14.4]	25.5 [18.3, NE]	12.9 [7.9, 18.5]
HR [95% CI]	0.59 [0.40, 0.89]		0.62 [0.43, 0.88]	
p-value	0.011		N/A	
<b>Secondary Endpoints - IRC-assessed</b>				
<b>Progression-Free Survival</b>				
Number (%) of patients with events	68 (37.2)	44 (48.4)	90 (49.2)	54 (59.3)
Median [95% CI], months	12.1 [6.8, 18.3]	3.3 [2.5, 5.6]	13.8 [8.7, 20.5]	3.6 [2.5, 7.1]
HR [95% CI]	0.37 [0.25, 0.55]		0.40 [0.28, 0.57]	
p-value	<0.001		N/A	
<b>Complete Response Rate</b>				
Responders (%)	92 (50.3)	20 (22.0)	107 (58.5)	23 (25.3)
Difference in response rate (%) [95% CI]	28.3 [16.3, 40.3]		33.2 [20.9, 45.5]	
p-value	<0.001		N/A	
<b>Objective Response Rate</b>				
Responders (%) (CR, PR)	110 (60.1)	29 (31.9)	125 (68.3)	37 (40.7)
Difference in response rate (%) [95% CI]	28.2 [15.5, 50.0]		27.7 [14.7, 40.6]	

CI=confidence interval; HR=hazard ratio; N/A=not applicable; NE=not estimable.

**Figure 3 Kaplan-Meier Plot of Overall Survival in Study GO41944 (STARGLO, Updated Analysis; ITT)**



**Figure 4 Kaplan-Meier Plot of IRC-Assessed Progression-Free Survival in Study GO41944 (STARGLO, Updated Analysis; ITT)**



**Patient-Reported Outcomes**

At baseline, patients in both treatment arms (Columvi+GemOx and R-GemOx) reported moderate to moderate-high levels of Physical Functioning and low levels of fatigue as measured by the EORTC QLQ-C30 and low to low-moderate levels of lymphoma symptoms as measured by the FACT-Lym LymS. During treatment, patients in both arms maintained their baseline level of physical functioning and experienced a similar deterioration in fatigue and lymphoma symptoms.

Median time to deterioration in the treatment arms in fatigue or physical functioning or lymphoma symptoms was comparable. Both arms experienced clinically meaningful improvement in pain, and Columvi+GemOx patients experienced larger improvements at earlier time points in pain as compared to R-GemOx patients. The results should be interpreted with caution in the context of the open-label study design.

**3.1.4 Immunogenicity**

As with all therapeutic proteins, there is a potential immunogenicity.

Across studies, the majority of patients (574/608 patients; 94.4%) who received Columvi were negative for antidrug antibodies (ADAs) at baseline and remained negative throughout treatment with Columvi. Four patients (0.7%) were negative for ADAs at baseline and became positive for ADAs during treatment. Four patients (0.7%) were ADA-positive at baseline and at one or more post-dose timepoints. Due to the limited number of patients with antibodies against glofitamab, no conclusions can be drawn concerning a potential effect of immunogenicity on efficacy or safety.

The detection of an immune response is highly dependent on the sensitivity and specificity of the assays used, sample handling, timing of sample collection, concomitant medications, and underlying disease. For these reasons, comparison of the incidence of antibodies to glofitamab with the incidence of antibodies to other products may be misleading.

**3.2 PHARMACOKINETIC PROPERTIES**

Non-compartmental analyses indicate that glofitamab serum concentration reaches the maximal level (C<sub>max</sub>) at the end of infusion and declines in a bi-exponential fashion. Glofitamab exhibits linear and dose-proportional pharmacokinetics over the dose range studied (0.005 to 30 mg) and is independent of time.

**3.2.1 Absorption**

Columvi is administered as an IV infusion. Peak concentration of glofitamab (C<sub>max</sub>) was reached at the end of the infusion.

**3.2.2 Distribution**

Following IV administration, the central volume of distribution was 3.34 L, which is close to total serum volume. The peripheral volume of distribution was 2.35 L.

**3.2.3 Metabolism**

The metabolism of glofitamab has not been directly studied. Antibodies are cleared principally by catabolism.

**3.2.4 Elimination**

The glofitamab serum concentration-time data are described by a population pharmacokinetic model with two compartments and both time-independent clearance and time-varying clearance.

The time-independent clearance pathway was estimated as 0.633 L/day and the initial time-varying clearance pathway as 0.814 L/day, with an exponential decay over time (K<sub>des</sub> ~ 1.5/day). The estimated decay half-life from the initial total clearance value to the time-independent clearance only was estimated as 0.471 days.

The effective half-life in the linear phase (i.e., after the contribution of time-varying clearance has collapsed to a negligible amount) is 7.92 days (geometric mean, 95% CI: 4.69, 11.90) based on the population pharmacokinetic analysis.

### 3.2.5 Pharmacokinetics in Special Populations

#### Pediatric Population

No studies have been conducted to investigate the pharmacokinetics of glofitamab in pediatric patients.

#### Geriatric Population

No differences in glofitamab exposure were noted in patients 65 years of age and older and those under 65 years based on population pharmacokinetic analysis.

#### Renal impairment

Population pharmacokinetic analyses showed that creatinine clearance does not affect the pharmacokinetics of glofitamab. The pharmacokinetics of glofitamab in patients with mild or moderate renal impairment (CrCL 30 to < 90 mL/min) were similar to those in patients with normal renal function. No dose adjustment is required for patients with mild or moderate renal impairment. Columvi has not been studied in patients with severe renal impairment.

#### Hepatic impairment

Population pharmacokinetic analyses showed hepatic impairment does not affect the pharmacokinetics of glofitamab. The pharmacokinetics of glofitamab in patients with mild hepatic impairment (total bilirubin > ULN to  $\leq 1.5 \times$  ULN or AST > ULN) were similar to those with normal hepatic functions. No dose adjustment is required for patients with mild hepatic impairment. Columvi has not been studied in patients with moderate and severe hepatic impairment.

## 3.3 NONCLINICAL SAFETY

### 3.3.1 Carcinogenicity

No carcinogenicity studies have been performed to establish the carcinogenic potential of Columvi.

### 3.3.2 Genotoxicity

No studies have been performed to establish the mutagenic potential of Columvi.

### 3.3.3 Impairment of Fertility

No fertility assessments in animals have been performed to evaluate the effect of Columvi.

### 3.3.4 Reproductive toxicity

No reproductive toxicity studies in animals have been performed to evaluate the effect of Columvi.

Based on low placental transfer of antibodies during the first trimester, the mechanism of action of glofitamab (B-cell depletion, target-dependent T-cell activation, and cytokine release), the available safety data with Columvi, and the data on other anti CD20 antibodies, the risk for teratogenicity is low. Prolonged B-cell depletion can lead to increased risk of opportunistic infection, which may cause fetal loss. Transient CRS associated with Columvi administration may also be harmful to the fetus.

### 3.3.5 Other

In a study in cynomolgus monkeys, animals experiencing severe CRS after a single intravenous dose of glofitamab (0.1 mg/kg) without obinutuzumab pre-treatment had erosions in the gastrointestinal tract and inflammatory cell infiltrates in spleen and sinusoids of the liver and sporadically in some other organs. These inflammatory cell infiltrates were likely secondary to cytokine-induced immune cell activation. Pre-treatment with obinutuzumab resulted in the attenuation of glofitamab-induced cytokine release and related adverse effects by depleting B cells in peripheral blood and lymphoid tissue. This allowed at least 10 times higher doses of glofitamab (1 mg/kg) in cynomolgus monkeys resulting in a  $C_{max}$  of up to 3.8 times the human  $C_{max}$  at the recommended 30 mg dose. All findings with glofitamab were considered pharmacologically mediated effects and reversible.

Studies longer than 4 weeks were not performed, as glofitamab was highly immunogenic in cynomolgus monkeys and led to loss of exposure and loss of the pharmacologic effect.

## 4. PHARMACEUTICAL PARTICULARS

### 4.1 STORAGE

#### Vials

Store at 2 °C to 8 °C.

Keep vial in the outer carton in order to protect from light.

Do not freeze. Do not shake.

#### Shelf life

As registered locally.

Columvi should not be used after the expiry date (EXP) shown on the carton.

#### Diluted solution for intravenous infusion

The prepared infusion solution should be used immediately. If not used immediately, the infusion solution can be stored in the refrigerator at 2 °C to 8 °C for up to 72 hours and at 30 °C for up to 24 hours, if prepared under aseptic conditions, followed by a maximum infusion time of 8 hours.

### 4.2 SPECIAL INSTRUCTIONS FOR USE, HANDLING AND DISPOSAL

Columvi diluted solution can be administered via intravenous bag infusion (all doses) or intravenous syringe infusion (2.5 mg dose only).

#### Instructions for dilutions

- Columvi contains no preservative and is intended for single use only.
- Columvi must be diluted by a healthcare professional using aseptic technique, prior to intravenous administration.
- Visually inspect the Columvi vial for particulate matter or discoloration prior to administration. Columvi is a colorless, clear solution. Discard the vial if the solution is cloudy, discolored, or contains visible particles.

#### Preparation of intravenous bag infusion

- Withdraw the required volume of 0.9% or 0.45% sodium chloride solution from the infusion bag (see Table 14) using a sterile needle and syringe and discard.
- Withdraw the required volume of Columvi concentrate for the intended dose from the vial using a sterile needle and syringe and dilute into the infusion bag (see Table 14). Discard any unused portion left in the vial.
- The final drug concentration after dilution must be 0.1 mg/mL to 0.6 mg/mL.
- Gently invert the infusion bag to mix the solution in order to avoid excessive foaming. Do not shake.
- Inspect the infusion bag for particulates and discard if present.
- Prior to the start of the intravenous infusion, the content of the infusion bag should be at room temperature.

Table 14 Dilution of Columvi for Intravenous Bag Infusion

Dose of Columvi to be administered	Size of 0.9% or 0.45% sodium chloride solution infusion bag	Volume of 0.9% or 0.45% sodium chloride solution to be withdrawn and discarded	Volume of Columvi concentrate to be added
2.5 mg	50 mL	27.5 mL	2.5 mL
	100 mL	77.5 mL	2.5 mL
10 mg	50 mL	10 mL	10 mL
	100 mL	10 mL	10 mL
30 mg	50 mL	30 mL	30 mL
	100 mL	30 mL	30 mL

#### Preparation of intravenous syringe infusion (2.5 mg dose only)

Use a two-syringe method with a connector to prepare the dose. The final volume of the diluted solution is 25 mL.

- Withdraw 22.5 mL of 0.9% or 0.45% sodium chloride solution from an infusion bag into a syringe of the appropriate size (e.g., 30 mL).
- Withdraw 2.5 mL of Columvi concentrate from the vial using a sterile needle into a second syringe. Discard any unused portion left in the vial.
- Attach a connector to the two syringes and transfer Columvi into the syringe containing 0.9% or 0.45% sodium chloride solution. The final drug concentration after dilution should be 0.1 mg/mL.
- Disconnect the syringes. Draw air into the syringe containing the Columvi diluted solution and close.

- Gently invert the syringe to mix the solution, in order to avoid excessive foaming. Do not shake.
- Remove air bubbles from the syringe before administration.

#### Administration

Only administer as an intravenous infusion.

Do not administer as an intravenous push or bolus.

Administer as an intravenous infusion through a dedicated infusion line using intravenous infusion pump or syringe pump, over a maximum of 8 hours.

Once the Columvi infusion bag or syringe is empty, ensure the entire dose of Columvi is administered by clearing the infusion line with an infusion bag or syringe containing 0.9% or 0.45% sodium chloride solution. Continue the infusion at the same rate until the recommended infusion duration is reached according to Table 2.

#### Incompatibilities

Only 0.9% or 0.45% sodium chloride solution should be used to dilute Columvi, since other diluents have not been tested.

Columvi when diluted with 0.9% sodium chloride solution is compatible with intravenous infusion bags composed of polyvinyl chloride (PVC), polyethylene (PE), polypropylene (PP), or non-PVC polyolefin. When diluted with 0.45% sodium chloride solution, Columvi is compatible with intravenous infusion bags composed of PVC.

Columvi when diluted with 0.9% or 0.45% sodium chloride solution is compatible with syringes composed of PP.

No incompatibilities have been observed with infusion sets with product-contacting surfaces of polyurethane (PUR), PVC, PE, polybutadiene (PBD), polyetherurethane (PEU), polycarbonate (PC), silicone, polytetrafluoroethylene (PTFE), or acrylonitrile butadiene styrene (ABS), and in-line filter membranes composed of polyethersulfone (PES) or polysulfone. The use of in-line filter membranes is optional.

#### Disposal of unused/expired medicines

The release of pharmaceuticals in the environment should be minimized. Medicines should not be disposed of via wastewater, and disposal through household waste should be avoided.

The following points should be strictly adhered to regarding the use and disposal of syringes and other medicinal sharps:

- Needles and syringes should never be reused.
- Place all used needles and syringes into a sharps container (puncture-proof disposable container).

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

#### Medicine: keep out of reach of children

Current at February 2026



F. Hoffmann-La Roche Ltd, Grenzachstrasse 124, CH-4070 Basel, Switzerland