

Article Summary

Cross-transmission in the dental office: Does this make you ill?

Original article by Volgenant & de Soet in
Current Oral Health Reports (2018) 5:221–228.

The article is a review paper of the literature relating to transmission of infections in dental practice. As the authors “aimed to provide the latest insights in the relative risks of transmission of (pathogenic) micro-organisms in the dental office” a number of new scientific publications is presented for which they discuss possible risks of transmission to patients through dental treatment. There is no method section outlining how the literature was searched, but a categorization is provided by the authors in the reference section to indicate important papers.

One major principle is stressed in the beginning: “The prevention of disease becomes more and more important in an era where increased antibiotic resistance results in a rise in untreatable infections.” A key element in disease prevention is understanding infection and transmission pathways. Accordingly, three modes of transmission enabling microorganisms to reach potential hosts are exemplified in the text.

Transmission by Direct contact: This can occur via hands, improper sterilised instruments or needle stick accidents. The magnitude of this type of cross-transmission is difficult to estimate as there are no active surveillance programmes in place for post-operative infections in dental practice. Examples given in this section are based on work undertaken on **Methacillin-resistant Staphylococcus aureus (MRSA)** carriage and transmission linked to dental staff or the dental environment. Further difficulties in estimating the risk of transmission in dental practices are highlighted since not all transmissions will result in infection. So, most of these will lead to transient colonization in the case of MRSA rather than infection. Additional studies are discussed, and the authors again showcase challenges in obtaining a good quality evidence base by indicating that the possibility for cross-transmission through dental equipment exists. Although undeniable proof is difficult to obtain, probably due to incomplete reporting and confounding variables.

Transmission by blood contact: A risk of transmission in the dental office exists when pathogens are transported directly from blood (e.g., of the patient) to blood (e.g., of the Dental Health Care Personnel (DHCP)). These blood exposure accidents (BEAs) are common in dentistry with frequent work with sharp instruments and needles. In a recent (2016) overview concerning transmission of blood-borne pathogens in the USA, Cleveland et al. found only three reports on cross-infection of HBV and HCV in the dental health care setting. Thus, it is concluded that there is a risk in the dental office, for transmission of blood-borne pathogens but this risk is low. From the perspective of health and safety for DHCP introduction of safety engineered devices or improved injection techniques to prevent BEAs have been recommended and should be adopted wherever possible in dental practice.

Transmission by dental unit water and aerosols: The most reported pathogens from contaminated water are Legionella and Pseudomonas species, but also, opportunistic genera such as Propioniumbacterium, Mycobacterium and Stenotrophomonas species are detected in the dental unit water line (DUWL). In dentistry two cases of legionellosis have been reported recently. However, despite the fact that two people died from a Legionella-pneumonia after a dental treatment, it still is debated whether (contaminated) DUWLs were the source of the Legionella, or that this bacterium had a different origin. Currently, no scientific evidence exists supporting an overall high occupational risk of Legionella infection from DUWL.

In discussion, it is suggested that the research reports described in the review underline the potential for transmission resulting in infection or carriage of micro-organisms. Consequently, maintaining a high standard of infection preventive measures for all patients must take a high priority. But the authors also emphasise their concerns that a number of studies conclude that the knowledge of DHCP about cross-transmission, cross-infection and how to prevent them is low.

A note on relative and absolute risks

A relative risk is a ratio of the probability of an event occurring in the exposed group (in this case patients exposed to dental treatment where there has been a deficiency in infection prevention) versus the probability of the event occurring in the non-exposed group (not receiving dental treatment). Relative risk does not provide any information about the absolute risk of the event occurring but rather the higher or lower likelihood of the event in the exposure versus the non-exposure group.

Thus, to calculate the relative risk, we must know the exposure status of all individuals (either exposed or not exposed). This implies that relative risk is only appropriate for cases where the exposure status and incidence of disease can be accurately determined such as prospective cohort studies – this is very difficult to undertake in dentistry unless one is investigating a specific exposure risk such as, BBV risk in a specific incident investigation.

Relative risks are often reported in newspaper headlines, but without the context of absolute (or baseline) risk, this information is meaningless. Absolute risk numbers are needed to understand the implications of relative risks and how specific factors or behaviours affect the likelihood of developing a disease or health condition.

Absolute risk is the actual risk of some event happening given the current exposure. Absolute risks are needed to understand relative risk. This is very difficult to calculate for routine dental treatment as there is little or no data on adverse events in treated and/or control groups.

For more details on risk see

<https://www.harding-center.mpg.de/en/persons/gerd-gigerenzer>

Or

D Spiegelhalter https://papers.ssrn.com/sol3/papers.cfm?abstract_id=2937730

Cited article:

Volgenant, C.M.C., de Soet, J.J. Cross-transmission in the Dental Office: Does This Make You Ill?. Curr Oral Health Rep 5, 221–228 (2018).
<https://doi.org/10.1007/s40496-018-0201-3>

Short notes on Transmission of infection

Prof AJ Smith

Background

How do infections occur?

An infection occurs when micro-organisms enter the body, increase in number, and cause a reaction of the body. **Three things are necessary for an infection to occur:**

- › Reservoir: Places where infectious agents live (e.g., sinks, surfaces, human skin)
- › Susceptible Person with a way for micro-organisms to enter the body
- › Transmission: a way micro-organisms are moved to the susceptible person

In healthcare settings, transmission of microbes depends on people, the environment and/or medical devices.

Transmission

The common mechanisms of transmission in healthcare settings, including dental practice are;

Through contaminated hands or gloves for example, **Methacillin-resistant Staphylococcus aureus (MRSA)** contamination on surfaces can be transmitted to patients or other staff via hands. Even whilst wearing latex gloves it is important that these are changed between patients. Many Gram positive bacteria such as MRSA can survive for months on dry surfaces. It has been reported that strains of Staphylococcus aureus can survive on dry surfaces for time periods between 7 days to 7 months (3). Review articles have also noted that bacteria survive on surfaces for longer if there are higher numbers of bacteria and the presence of proteins included as such as serum, sputum or dust (5). Similar MRSA strains were recovered in patients

and dental surgery samples after attending a dental clinic (4) indicating transmission from clinic surfaces. Strains of *Staphylococcus aureus* can be recovered from the surfaces of portable electronic devices such as computers and ipads (Khan et al AJIC 2015). Little work has been undertaken investigating recovery of respiratory tract viruses from dental surfaces, although there has been an investigation of the immune response in dentists to assess their exposure to respiratory virus infections (Davies et al BDJ 1994 176: 262-5) that showed general dental practitioners had significantly raised antibody titres compared to controls for influenza A, B and respiratory syncytial virus.

1

Sprays or splashes from dental aerosols, can spread upper respiratory tract viruses, such as influenza. Large contaminated droplets can contaminate surfaces and transmission via hands. Influenza A and B can survive on steel and plastic surfaces for 24–48 hours and cloth, paper and tissues for <8-12 hours (Bean et al. J Infect Dis 2002). Transmission can occur for influenza viruses to steel to hands over a 24 hour period and the virus can survive for approximately 5 minutes on the hands (1). Smaller aerosolised particles can be inhaled or contaminate eyes to cause infection. Attention must also be paid to following manufacturer's instructions on the disinfection of dental unit waterlines.

2

Sharp injuries can lead to transmission of infection, for example, Hepatitis B virus can be transmitted when the skin is punctured by a used needle. The risk of Hepatitis B transmission through a contaminated sharps injury from a Hepatitis B positive patient to a non-immune recipient is estimated at 30% (2). A relatively recent example of transmission of Hepatitis B through dental procedures that demonstrates its potentially high infectious nature is that associated with a portable dental clinic in the USA where three patients and two staff were infected with the Hepatitis B virus (Radcliffe et al. J Am Dent Assoc. 2013;144(10):1110-8.). The precise mode of transmission was not detected but multiple breaches in protocols were noted such as close proximity of clean and contaminated instruments, dental handpieces were not sterilized between patients and there was no written records or traceability for the processes used.

What determines the rate of transmission?

The basic reproduction ratio (R_0) is a measure of the ability of a pathogen to give rise to more infections or secondary cases. The rate of transmission of a disease is determined by a number of factors in the infectious agent.

Such as, virulence (ability to cause disease) for example the pandemic influenza strains from 1918 is reported to have a higher R_0 compared to other flu viruses and host factors, such as immune response (vaccines) – the influenza vaccine is predicted each year to try and match against circulating flu viruses each year. Therefore, the efficacy of the virus can vary from season to season. Nevertheless, it is essential that dental healthcare workers get vaccinated against circulating flu viruses each year to protect themselves and vulnerable patients they may be treating.

If the R_0 is greater than 1 then the infection has potential to spread through a susceptible population. The higher the number the higher the rate of transmission. For measles, R_0 is often cited to be 12–18, which means that each person with measles would, on average, infect 12–18 other people in a totally susceptible population. The **R_0** for novel **influenza A (H1N1)** has recently been estimated to be between 1.4 and 1.6. It has been estimated the basic reproduction number, R_0 to be 1.53 for Hepatitis B (in New Zealand), and shown that the vaccination campaign against Hepatitis B has substantially reduced this below one. R_0 estimates for community strains of MRSA (USA 300) the R_0 ranged from 1.24 to 1.34.

Protection against transmission

The most practical method to prevent cross-transmission in dental practice is to adhere to standard infection control precautions for all patients as it is difficult to determine which microorganisms patients maybe carrying. It is also vital that all dental staff have received the recommend vaccines and that these are kept up to date and recorded in the practice record management system.

Bibliography:

- (1) Coburn BJ, Wagner BG, Blower S. Modeling influenza epidemics and pandemics: insights into the future of swine flu (H1N1). BMC Med. 2009 Jun 22;7:30
- (2) Mann J, Roberts M. Modelling the epidemiology of hepatitis B in New Zealand. J Theor Biol. 2011 Jan 21;269(1):266-72.
- (3) Prosperi M et al., Molecular Epidemiology of Community-Associated Methicillin-resistant Staphylococcus aureus in the genomic era: a Cross-Sectional Study. Sci Rep 2013; 3, 1902
- (4) Kurita et al. Nosocomial transmission of MRSA via the surfaces of the dental operator. BDJ 2006
- (5) Kramer et al, BMC Infectious Diseases 2006